

2

AD-A242 589



DOCUMENTATION PAGE

Form Approved  
OMB No 0704-0188

18 1991

1b. RESTRICTIVE MARKINGS

3. DISTRIBUTION / AVAILABILITY OF REPORT

Approval for public release  
Distribution list enclosed

2b. DECLASSIFICATION / DOWNGRADING SCHEDULE

4. PERFORMING ORGANIZATION REPORT NUMBER(S)

Technical Report No. 30

5. MONITORING ORGANIZATION REPORT NUMBER(S)

6a. NAME OF PERFORMING ORGANIZATION  
State University of New York  
at Buffalo6b. OFFICE SYMBOL  
(If applicable)7a. NAME OF MONITORING ORGANIZATION  
Office of Naval Research6c. ADDRESS (City, State, and ZIP Code)  
Department of Chemistry  
Buffalo, NY 142147b. ADDRESS (City, State, and ZIP Code)  
Department of the Navy  
800 North Quincy Street  
Arlington, VA 22217-50008a. NAME OF FUNDING / SPONSORING  
ORGANIZATION  
Office of Naval Research8b. OFFICE SYMBOL  
(If applicable)9. PROCUREMENT INSTRUMENT IDENTIFICATION NUMBER  
N00014-90-J-1530; R&T Code 41350028c. ADDRESS (City, State, and ZIP Code)  
Department of the Navy  
Arlington, VA 22217

10. SOURCE OF FUNDING NUMBERS

PROGRAM  
ELEMENT NOPROJECT  
NO.TASK  
NOWORK UNIT  
ACCESSION NO  
4135002

11. TITLE (Include Security Classification)

Organogallium Compounds Containing the Neophyl Substituent. The Crystal and Molecular Structure of trans-[(PhMe<sub>2</sub>CCH<sub>2</sub>)<sub>2</sub>GaN(H)(Pr)]<sub>2</sub>.

12. PERSONAL AUTHOR(S)

O. T. Beachley, Jr.\*, M. J. Noble, Melvyn Rowen Churchill\* and Charles H. Lake

13a. TYPE OF REPORT

Technical Report

13b. TIME COVERED

FROM \_\_\_\_\_ TO \_\_\_\_\_

14. DATE OF REPORT (Year, Month, Day)

11 November 1991

15. PAGE COUNT

27

16. SUPPLEMENTARY NOTATION

To be published in Organometallics

17. COSATI CODES

| FIELD | GROUP | SUB-GROUP |
|-------|-------|-----------|
|       |       |           |
|       |       |           |
|       |       |           |

18. SUBJECT TERMS (Continue on reverse if necessary and identify by block number)

Gallium chemistry, organogallium compounds, gallium-nitrogen compounds, x-ray structural study

19. ABSTRACT (Continue on reverse if necessary and identify by block number)

A series of organogallium compounds which contain the neophyl substituent including Ga(CH<sub>2</sub>CMe<sub>2</sub>Ph)<sub>3</sub>, Ga(CH<sub>2</sub>CMe<sub>2</sub>Ph)<sub>2</sub>Cl<sub>2</sub> and (PhMe<sub>2</sub>CCH<sub>2</sub>)<sub>2</sub>GaN(H)(Pr) has been prepared in high yield by reactions typical of organogallium chemistry. The characterization data include elemental analyses, melting points, NMR and IR spectroscopic studies and cryoscopic molecular weight studies. All data are consistent with the conclusions that Ga(CH<sub>2</sub>CMe<sub>2</sub>Ph)<sub>3</sub> is monomeric in benzene solution whereas Ga(CH<sub>2</sub>CMe<sub>2</sub>Ph)<sub>2</sub>Cl<sub>2</sub>, Ga(CH<sub>2</sub>CMe<sub>2</sub>Ph)Cl<sub>2</sub> and (PhMe<sub>2</sub>CCH<sub>2</sub>)<sub>2</sub>GaN(H)(Pr) are dimeric. The gallium amide [(PhMe<sub>2</sub>CCH<sub>2</sub>)<sub>2</sub>GaN(H)(Pr)]<sub>2</sub> crystallizes in the centrosymmetric triclinic space group P1̄ (No. 2) with a = 10.739(2), b = 10.844(2), c = 11.125(2) Å, α = 69.14(1), β = 62.96(1), γ = 76.74(1)°, V = 1074.7(3) Å<sup>3</sup> and Z = 1. Diffraction data (MoKα, 2θ = 5-50°) were collected on a Siemens R3m/V automated four-circle diffractometer and the structure was refined to R = 3.57% and wR = 3.46% for all 3803 unique reflections (R = 2.52%, wR = 2.62% for those 3111 data with |F<sub>o</sub>| > 6σ(|F<sub>o</sub>|)). The molecule lies on a crystallographic inversion center and thus has a strictly planar Ga<sub>2</sub>N<sub>2</sub> core. Distances of interest include Ga-CH<sub>2</sub>CMe<sub>2</sub>Ph = 1.992(3) Å and Ga-N(bridging) = 2.013(2) and 2.029(2) Å.

20. DISTRIBUTION / AVAILABILITY OF ABSTRACT

☒ UNCLASSIFIED/UNLIMITED ☐ SAME AS RPT ☐ DTIC USERS

21. ABSTRACT SECURITY CLASSIFICATION

Unclassified

22a. NAME OF RESPONSIBLE INDIVIDUAL

O. T. Beachley, Jr.

22b. TELEPHONE (Include Area Code)

(716) 831-3266

22c. OFFICE SYMBOL

DD Form 1473, JUN 86

Previous editions are obsolete

SECURITY CLASSIFICATION OF THIS PAGE

This document has been approved  
for public release and sale; its  
distribution is unlimited.

Unclassified

OFFICE OF NAVAL RESEARCH  
Contract N-00014-90-J-1530

R&T Code 4135002

TECHNICAL REPORT NO. 30

Organogallium Compounds Containing the Neophyl Substituent.  
The Crystal and Molecular Structure of trans-[(PhMe<sub>2</sub>CCH<sub>2</sub>)<sub>2</sub>GaN(H)(Pr)]<sub>2</sub>

by

O. T. Beachley, Jr.<sup>\*</sup>, M. J. Noble, Melvyn Rowen Churchill<sup>\*</sup>  
and Charles H. Lake

Prepared for Publication  
in  
Organometallics

State University of New York at Buffalo  
Department of Chemistry  
State University of New York at Buffalo, Buffalo, NY 14214

11 November 1991

Reproduction in whole or in part is permitted for  
any purpose of the United States Government

<sup>\*</sup>This document has been approved for public release  
and sale; its distribution is unlimited

01 1115 059

91-15753



Organogallium Compounds Containing the Neophyl Substituent.  
The Crystal and Molecular Structure of trans-[(PhMe<sub>2</sub>CCH<sub>2</sub>)<sub>2</sub>GaN(H)(Pr)]<sub>2</sub>

by

O. T. Beachley, Jr.<sup>\*</sup>, M. J. Noble, Melvyn Rowen Churchill<sup>\*</sup>

and Charles H. Lake

Department of Chemistry  
State University of New York at Buffalo, Buffalo, NY 14214

Abstract

A series of organogallium compounds which contain the neophyl substituent including Ga(CH<sub>2</sub>CMe<sub>2</sub>Ph)<sub>3</sub>, Ga(CH<sub>2</sub>CMe<sub>2</sub>Ph)<sub>2</sub>Cl, Ga(CH<sub>2</sub>CMe<sub>2</sub>Ph)Cl<sub>2</sub> and (PhMe<sub>2</sub>CCH<sub>2</sub>)<sub>2</sub>GaN(H)(Pr) has been prepared in high yield by reactions typical of organogallium chemistry. The characterization data include elemental analyses, melting points, NMR and IR spectroscopic studies and cryoscopic molecular weight studies. All data are consistent with the conclusions that Ga(CH<sub>2</sub>CMe<sub>2</sub>Ph)<sub>3</sub> is monomeric in benzene solution whereas Ga(CH<sub>2</sub>CMe<sub>2</sub>Ph)<sub>2</sub>Cl, Ga(CH<sub>2</sub>CMe<sub>2</sub>Ph)Cl<sub>2</sub> and (PhMe<sub>2</sub>CCH<sub>2</sub>)<sub>2</sub>GaN(H)(Pr) are dimeric. The gallium amide [(PhMe<sub>2</sub>CCH<sub>2</sub>)<sub>2</sub>GaN(H)(Pr)]<sub>2</sub> crystallizes in the centrosymmetric triclinic space group P $\bar{1}$  (No. 2) with  $a = 10.739(2)$ ,  $b = 10.844(2)$ ,  $c = 11.125(2)$  Å,  $\alpha = 69.14(1)$ ,  $\beta = 62.96(1)$ ,  $\gamma = 76.74(1)^\circ$ ,  $V = 1074.7(3)$  Å<sup>3</sup> and  $Z = 1$ . Diffraction data (Mo K $\alpha$ ,  $2\theta = 5-50^\circ$ ) were collected on a Siemens R3m/V automated four-circle diffractometer and the structure was refined to  $R = 3.57\%$  and  $wR = 3.46\%$  for all 3803 unique reflections ( $R = 2.52\%$ ,  $wR = 2.62\%$

A-1

for those 3111 data with  $|F_o| > 6\sigma(|F_o|)$ . The molecule lies on a crystallographic inversion center and thus has a strictly planar  $\text{Ga}_2\text{N}_2$  core. Distances of interest include  $\text{Ga}-\text{CH}_2\text{CMe}_2\text{Ph} = 1.992(2)$  and  $1.994(3)\text{\AA}$  and  $\text{Ga}-\text{N}(\text{bridging}) = 2.013(2)$  and  $2.029(2)\text{\AA}$ .

### Introduction

The utilization of gallium in electronic materials including semiconductors and superconductors has led to a renewed interest in its chemistry. Even though a variety of organogallium compounds are known, relatively little is known about compounds which incorporate bulky organic substituents. Bulky substituents can lead to compounds with enhanced thermal stability as well as to compounds with unexpected structures. Consequently, our prior research involved investigations of the chemistry of gallium compounds with  $\text{CH}_2\text{SiMe}_3$ ,<sup>1</sup>  $\text{CH}_2\text{CMe}_3$ ,<sup>2</sup>  $2,4,6\text{-Me}_3\text{-C}_6\text{H}_2$ ,<sup>3,4</sup> and  $\text{C}_5\text{Me}_5$ ,<sup>5,6</sup> substituents but only the mesityl derivative  $\text{Ga}(\text{C}_6\text{Me}_3\text{H}_2)\text{Cl}_2$  displayed unusual chemistry. The compound  $\text{Ga}(\text{C}_6\text{Me}_3\text{H}_2)\text{Cl}_2$  was characterized as a unique one dimensional polymer<sup>4</sup> in which planar  $\text{Ga}(\text{C}_6\text{Me}_3\text{H}_2)\text{Cl}$  units with distorted tetrahedral geometry were linked by bridging chloride ligands. The traditional structure of an organogallium dichloride is a chloride bridged dimer.

In this paper the synthesis and characterization of organogallium compounds which incorporate the bulky neophyl ligand including  $\text{Ga}(\text{CH}_2\text{CMe}_2\text{Ph})_3$ ,  $\text{Ga}(\text{CH}_2\text{CMe}_2\text{Ph})_2\text{Cl}$ ,  $\text{Ga}(\text{CH}_2\text{CMe}_2\text{Ph})\text{Cl}_2$  and  $(\text{PhMe}_2\text{CCH}_2)_2\text{GaN}(\text{H})(\text{Pr})$  are described.

### EXPERIMENTAL

All compounds described in this investigation were very sensitive to oxygen and moisture and were manipulated by standard vacuum line techniques or under a purified argon atmosphere. All solvents were purified before use. Gallium(III) chloride was purified by sublimation under dynamic high vacuum at 80-85 °C immediately prior to use. Neophyl chloride was purchased from Aldrich Chemical Co. and was distilled prior to use. Elemental

analyses were performed by Schwarzkopf Microanalytical Laboratory, Woodside, New York or by E+R Microanalytical Laboratories, Corona, New York. Infrared spectra of Nujol mulls or of the neat compound between CsI plates were recorded by means of a Perkin-Elmer 683 Spectrometer. Absorption intensities are reported with the abbreviations vs (very strong), s (strong), m (medium), w (weak), vw (very weak), sh (shoulder). The  $^1\text{H}$  NMR spectra were recorded at 300 MHz or 90 MHz by using either a Varian Gemini 300 or Varian EM-390 spectrometer, respectively. All samples for NMR spectra were contained in sealed NMR tubes. Chemical shifts are reported in  $\delta$  units (ppm) and are referenced to tetramethylsilane (TMS) as  $\delta = 0.00$  ppm and benzene as  $\delta = 7.15$  ppm. Melting points were observed in a Mel-Temp by using sealed capillaries. Molecular weights were measured cryoscopically in benzene by using an instrument similar to that described by Shriver and Dreznor.<sup>7</sup>

Synthesis of  $\text{Ga}(\text{CH}_2\text{CMe}_2\text{Ph})_3$ . In a typical experiment the Grignard reagent,  $(\text{PhMe}_2\text{CCH}_2)\text{MgCl}$ , was prepared from magnesium turnings (6.217 g, 255.8 mmol) and freshly distilled neophyl chloride (41.193 g, 244.22 mmol) in 100 mL of diethyl ether. The magnesium turnings were activated with iodine without stirring prior to addition of the neophyl chloride/ether solution. The reaction mixture was stirred under reflux for 18 h. A flask charged with 10.0 g (56.9 mmol) of  $\text{GaCl}_3$  dissolved in 250 mL of dry diethyl ether was fitted with a condenser, mechanical stirrer, and a pressure equalizing addition funnel. Under a cover of argon the Grignard reagent solution was transferred to the addition funnel and was added to the  $\text{GaCl}_3$  solution over a period of 30 minutes. After the addition was complete, the reaction mixture was stirred under reflux for 18 h. The  $\text{Et}_2\text{O}$  was then removed by vacuum distillation at room temperature. The crude product was

extracted from the  $\text{MgCl}_2$  by 4-5 washings with 150 mL of pentane to provide a yellowish "wet" solid which was purified by vacuum distillation in a short-path still at 120-210 °C. The first fraction which distilled at 120-150 °C was identified as bineophyl  $\text{PhMe}_2\text{CCH}_2\text{CH}_2\text{CMe}_2\text{Ph}$  by its  $^1\text{H}$  NMR spectrum<sup>8,9</sup> and mp.<sup>8,9</sup> The second fraction which distilled at 160-210 °C was identified as pure  $\text{Ga}(\text{CH}_2\text{CMe}_2\text{Ph})_3$  and was obtained as a colorless solid (19.8 g, 42.2 mmol, 74.1% yield based on  $\text{GaCl}_3$ ). More than one distillation was frequently required for complete removal of the bineophyl. The product isolated by distillation can be further purified, if needed, by recrystallization from pentane at -20 °C.

$\text{Ga}(\text{CH}_2\text{CMe}_2\text{Ph})_3$ . mp 37.9-39.2 °C.  $^1\text{H}$  NMR ( $\text{C}_6\text{D}_6$ ,  $\delta$ ): 7.08 (m, 15H,  $-\text{C}_6\text{H}_5$ ); 1.23 (s, 18H,  $-\text{CMe}_2$ ), 0.81 (s, 6H,  $-\text{CH}_2-$ ). IR (Nujol mull,  $\text{cm}^{-1}$ ): 3110(sh, m), 3085(s), 3059(s), 3035(s), 3021(s), 1960(sh, w), 1941(w), 1885(w), 1869(w), 1800(w), 1742(w), 1668(w), 1598(s), 1579(m), 1535(w), 1490(vs), 1405(m), 1305(w), 1273(m), 1181(s), 1155(m), 1098(m), 1070(s), 1065(sh, m), 1028(vs), 998(w), 961(w), 925(w), 901(m), 858(w), 838(w), 761(vs), 725(sh, s), 698(vs), 660(sh, m), 620(m), 612(m), 551(s), 505(w), 457(w), 401(w), 269(w). Anal. calcd.: C, 76.77; H, 8.38. Found: C, 76.76; H, 8.56. Cryoscopic molecular weight, benzene solution, formula weight 469.36 (obsd. molality, obsd. mol. wt., association): 0.0846, 500.1, 1.07; 0.0536, 516.6, 1.10; 0.0372, 548.1, 1.17. Solubility: soluble in THF,  $\text{Et}_2\text{O}$ , benzene, and pentane. Trisneophylgallium(III) formed a stable 1:1 adduct with  $\text{NMe}_3$ , formed a weak 1:1 adduct with THF that slowly dissociated under dynamic vacuum, but did not form a stable adduct with  $\text{Et}_2\text{O}$ .

Synthesis of  $(\text{PhMe}_2\text{CCH}_2)_2\text{GaCl}$ . In a typical experiment,  $\text{Ga}(\text{CH}_2\text{CMe}_2\text{Ph})_3$  (2.854 g, 6.081 mmol) and  $\text{GaCl}_3$  (0.5353 g, 3.040 mmol) were weighed into

vials in the dry box. Both compounds were dissolved in pentane and then were pipetted into a 100 mL side arm flask. Each vial was rinsed several times (5-7) with fresh aliquots of pentane to ensure the quantitative transfer of reagents. After stirring the mixture at room temperature for 24 hours, the slightly cloudy solution was filtered through a medium glass frit. The product was then recrystallized from pentane at  $-10\text{ }^{\circ}\text{C}$  to yield pure  $(\text{PhMe}_2\text{CCH}_2)_2\text{GaCl}$  as a colorless solid (3.135 g, 8.436 mmol, 92.5% yield).

$(\text{PhMe}_2\text{CCH}_2)_2\text{GaCl}$ . mp  $86.4\text{--}87.8\text{ }^{\circ}\text{C}$ .  $^1\text{H}$  NMR ( $\text{C}_6\text{D}_6$ ,  $\delta$ ): 7.15 (m, 10H,  $-\text{C}_6\text{H}_5$ ); 1.35 (s, 4H,  $-\text{CH}_2-$ ); 1.28 (s, 12H,  $-\text{CMe}_2$ ). IR (Nujol mull,  $\text{cm}^{-1}$ ): 3082(s), 3055(s), 3019(vs), 2722(s), 1965(m), 1954(m), 1898(m), 1880(m), 1867(m), 1809(m), 1799(m), 1755(m), 1740(m), 1595(s), 1576(m), 1490(vs), 1398(sh, m), 1272(s), 1189(m), 1181(s), 1153(m), 1141(m), 1115(m), 1101(m), 1070(s), 1065(sh, m), 1025(s), 1001(m), 995(m), 948(m), 932(m), 928(m), 910(m), 901(m), 855(m), 835(sh, w), 768(sh, s), 761(s), 733(s), 719(s), 695(vs), 652(m), 612(w), 555(m), 512(w), 435(w), 420(w), 402(w), 315(vs), 280(w), 258(m), 242(sh, m). Anal. calcd.: C, 64.64; H, 7.05. Found: C, 64.35; H, 7.05. Cryoscopic molecular weight, benzene solution, formula weight 371.60 (obsd. molality, obsd. mol. wt., association): 0.0775, 773.2, 2.08; 0.0504, 769.4, 2.07; 0.0402, 769.1, 2.07. Solubility: soluble in THF,  $\text{Et}_2\text{O}$ , benzene, and pentane. Bisneophylgallium(III) chloride formed a stable 1:1 adduct with  $\text{NMe}_3$  but did not form a stable adduct with  $\text{Et}_2\text{O}$  or THF.

Synthesis of  $(\text{PhMe}_2\text{CCH}_2)_2\text{GaCl}_2$ . Neophylgallium(III) dichloride,  $(\text{CH}_2\text{CMe}_2\text{Ph})\text{GaCl}_2$ , was prepared by a stoichiometric redistribution reaction in the same manner as previously described for  $(\text{PhMe}_2\text{CCH}_2)_2\text{GaCl}$ . In a typical experiment  $\text{Ga}(\text{CH}_2\text{CMe}_2\text{Ph})_3$  (3.042 g, 6.480 mmol) and  $\text{GaCl}_3$  (2.282 g,



12.96 mmol) were reacted in pentane to yield  $\text{Ga}(\text{CH}_2\text{CMe}_2\text{Ph})\text{Cl}_2$  (4.849 g, 17.71 mmol, 91.1%). The compound was purified by recrystallization from pentane at  $-30\text{ }^\circ\text{C}$ .

$\text{Ga}(\text{CH}_2\text{CMe}_2\text{Ph})\text{Cl}_2$ . mp  $69.5\text{--}70.8\text{ }^\circ\text{C}$ .  $^1\text{H}$  NMR ( $\text{C}_6\text{D}_6$ ,  $\delta$ ): 7.14 (m, 5H,  $-\text{C}_6\text{H}_5$ ), 1.52 (s, 2H,  $-\text{CH}_2-$ ), 1.23 (s, 6H,  $-\text{CMe}_2$ ). IR (Nujol mull,  $\text{cm}^{-1}$ ): 3110(w), 3090(m), 3059(vs), 3025(vs), 2342(vw), 2319(w), 1965(m), 1948(m), 1890(m), 1871(m), 1820(w), 1800(m), 1778(w), 1746(m), 1680(w), 1598(s), 1579(m), 1539(m), 1509(m), 1490(vs), 1402(sh, m), 1395(s), 1309(m), 1288(sh, m), 1272(s), 1240(sh, m), 1205(sh, m), 1190(s), 1155(m), 1131(m), 1112(m), 1102(m), 1075(m), 1062(s), 1028(s), 1005(m), 995(m), 980(m), 960(m), 945(m), 931(m), 905(m), 862(m), 835(m), 767(vs), 729(vs), 705(sh, s), 694(vs), 639(s), 554(s), 509(m), 440(m), 415(m), 380(vs), 350(s), 260(vs), 230(vs). Anal. calcd.: C, 43.86; H, 4.79. Found: C, 43.92; H, 4.96. Cryoscopic molecular weight, benzene solution, formula weight 273.83 (obsd. molality, obsd. mol. wt., association): 0.0831, 588.5, 2.15; 0.0574, 569.7, 2.08; 0.0428, 573.2, 2.09. Solubility: soluble in THF,  $\text{Et}_2\text{O}$ , benzene, and pentane. Neophylgallium(III) dichloride formed a stable 1:1 adduct with  $\text{NMe}_3$  and THF but did not form a stable adduct with  $\text{Et}_2\text{O}$ .

Synthesis of  $(\text{PhMe}_2\text{CCH}_2)_2\text{GaN}(\text{H})(\text{Pr})$ . In a typical experiment,  $\text{Ga}(\text{CH}_2\text{CMe}_2\text{Ph})_3$  (2.16 g, 4.60 mmol) and propylamine (0.314 g, 5.31 mmol) were heated at  $140\text{--}150\text{ }^\circ\text{C}$  for 3d in a sealed tube. After heating, the volatile product from the elimination reaction  $\text{C}_6\text{H}_5\text{CMe}_3$  and excess  $\text{NH}_2\text{Pr}$  were removed by vacuum distillation. The organogallium amide  $(\text{PhMe}_2\text{CCH}_2)_2\text{GaN}(\text{H})\text{Pr}$  was isolated by washing the product out of the reaction tube with pentane through a medium glass frit and was recrystallized at  $-30\text{ }^\circ\text{C}$  to yield 1.514 g, 3.840 mmol, 83.4% yield based on  $\text{Ga}(\text{CH}_2\text{CMe}_2\text{Ph})_3$ . Crystallographic

quality crystals were obtained by slow cooling of a saturated pentane solution to -30 °C.

(PhMe<sub>2</sub>CCH<sub>2</sub>)<sub>2</sub>GaN(H)(Pr). mp 129.3-131.0 °C. <sup>1</sup>H NMR (C<sub>6</sub>D<sub>6</sub>, δ): 7.37 (d, 4H, o-Ar, <sup>3</sup>J<sub>CCH</sub> = 7.50 hz); 7.21 (t, 4H, m-Ar, <sup>3</sup>J<sub>CCH</sub> = 7.50 hz); 7.06 (t, 2H, p-Ar, <sup>3</sup>J<sub>CCH</sub> = 7.20 hz); 2.11 (q, 2H, -CH<sub>2</sub>-, <sup>3</sup>J<sub>CCH</sub> = 7.90 hz); 1.38 (s, 12H, -CMe<sub>2</sub>(neophyl)); 0.92 (s, 4H, -CH<sub>2</sub>(neophyl)-); 0.87 (m, 2H, -NCH<sub>2</sub>-); 0.66 (t, 3H, -CH<sub>3</sub>, <sup>3</sup>J<sub>CCH</sub> = 7.05 hz); 0.02 (t, 1H, -NH, <sup>3</sup>J<sub>CNH</sub> = 7.65 hz). IR (Nujol mull, cm<sup>-1</sup>): 3298(s), 3082(s), 3050(s), 3019(vs), 2720(s), 1955(m), 1940(m), 1595(s), 1578(m), 1531(m), 1490(vs), 1439(sh, vs), 1361(vs), 1356(s), 1301(m), 1276(m), 1245(m), 1230(m), 1225(m), 1181(s), 1154(m), 1130(m), 1098(w), 1085(m), 1071(m), 1061(sh, m), 1052(vs), 1045(sh, s), 1025(s), 1010(s), 999(m), 989(sh, m), 979(m), 955(m), 926(w), 900(m), 889(m), 862(s), 845(s), 829(sh, m), 760(vs), 748(m), 715(vs), 695(vs), 686(s), 620(m), 599(m), 568(s), 550(m), 505(w), 499(m), 459(s), 439(sh, w), 409(sh, w), 399(m), 268(vw), 236(m). Anal. Calcd.: C, 70.07; H, 8.69. Found: C, 70.24; H, 8.75. Cryoscopic molecular weight, benzene solution, formula weight 394.25 (obsd. molality, obsd. mol. wt., association): 0.0833, 709, 1.80; 0.0663, 713, 1.81; 0.0508, 723, 1.83. Solubility: soluble in benzene and pentane.

Collection of X-Ray Diffraction Data for [(PhMe<sub>2</sub>CCH<sub>2</sub>)<sub>2</sub>GaN(H)(Pr)]<sub>2</sub>.

The crystal selected for the structural study had well defined faces and approximate dimensions of 0.35mm × 0.30mm × 0.30mm. Due to its extreme air sensitivity it was sealed into a thin walled capillary under strictly anaerobic conditions. The capillary was then mounted and aligned on a Siemens R3m/V four-circle single-crystal diffractometer. Details of data collection appear in Table 1.

The crystal belongs to the triclinic system, possible space groups being the centrosymmetric  $P\bar{1}$  and the noncentrosymmetric  $P1$ . (The former was shown later to be the correct choice). An entire sphere of data (two equivalent forms) was collected for  $2\theta = 5-50^\circ$  (Mo K $\alpha$  radiation). This was done to increase the quality of the crystal structure through the merging of the two forms of data. A total of 6274 reflections were collected and merged to 3803 unique data ( $R_{\text{int}} = 1.8\%$ ). The resulting data were corrected for Lorentz and polarization effects. Absorption problems were corrected for by the use of a semi-empirical  $\psi$ -scan technique.

Solution and Refinement of the Structure of  $[(\text{PhMe}_2\text{CCH}_2)_2\text{Ga}(\text{H})(\text{Pr})]_2$ .

The resulting data set was fed into a VAX3100 workstation which was operating with the Siemens SHELXTL PLUS program system.<sup>10</sup> Data reduction, structural solution and data refinement were performed under this system.

The resulting solution was in the centrosymmetric spacegroup  $P\bar{1}$  ( $C_i^1$ ; No. 2). The structure was solved by the use of direct methods followed by full matrix least-squares refinement. Positional and anisotropic thermal parameters for all non-hydrogen atoms and isotropic thermal parameters for the hydrogen atoms were refined as well as positional parameters for the unique N-bonded hydrogen atom of the  $\mu$ -NHPr ligand. All remaining hydrogen atoms were input in calculated positions with  $d(\text{C-H}) = 0.96\text{\AA}$ .<sup>11</sup> Following refinement, the largest features remaining on the electron density difference map were a maximum peak height of  $0.30 \text{ e}^-/\text{\AA}^3$  and a minimum of  $-0.22 \text{ e}^-/\text{\AA}^3$ . This corresponded to a structural model which converged with reliability factors of  $R = 3.57\%$  and  $wR = 3.16\%$  for all 3803 unique data. ( $R = 2.52\%$  and  $wR = 2.62\%$  for those 3111 data with  $|F_o| > 6\sigma(|F_o|)$ .) The structure was also corrected for secondary extinction by refining the

parameter  $\chi$  in the expression  $F^* = F[1+0.002\chi F^2/\sin(w\theta)]^{-1/4}$ ; the refined value was  $\chi = 0.0004(2)$ .

Final atomic coordinates are collected in Table 2.

Lewis Acid-Base Studies. The tendency of  $\text{Ga}(\text{CH}_2\text{CMe}_2\text{Ph})_3$ ,  $(\text{PhMe}_2\text{CCH}_2)_2\text{GaCl}$ , and  $(\text{PhMe}_2\text{CCH}_2)_2\text{GaCl}_2$  to form Lewis acid-base adducts with  $\text{NMe}_3$ ,  $\text{Et}_2\text{O}$  and THF was investigated. In a typical experiment a weighed amount of excess Lewis base was vacuum distilled onto a weighed amount of the Lewis acid. After the resulting sample was warmed to room temperature and stirred for 1-2 hours, the excess base was removed by vacuum distillation. Mass measurements were used to determine the quantity of base which reacted. The mole ratio of Lewis acid to base which reacted, the melting point and the  $^1\text{H}$  NMR spectrum of the resulting adducts are described below. The Lewis acids  $\text{Ga}(\text{CH}_2\text{CMe}_2\text{Ph})_3$ ,  $(\text{PhMe}_2\text{CCH}_2)_2\text{GaCl}$  and  $(\text{PhMe}_2\text{CCH}_2)\text{GaCl}_2$  did not form stable adducts with  $\text{Et}_2\text{O}$  or  $(\text{PhMe}_2\text{CCH}_2)_2\text{GaCl}$  with THF.

$(\text{PhMe}_2\text{CCH}_2)_3\text{Ga}\cdot\text{NMe}_3$ . Mol ratio  $\text{Ga}(\text{CH}_2\text{CMe}_2\text{Ph})_3/\text{NMe}_3$ : 0.99. mp 119-134 °C (dec.).  $^1\text{H}$  NMR ( $\text{C}_6\text{H}_6$ ,  $\delta$ ): 1.61 (s, 9H,  $\text{NMe}_3$ ), 1.38 (s, 18H,  $-\text{CMe}_2$ ), 0.73 (s, 6H,  $-\text{CH}_2-$ ).

$(\text{PhMe}_2\text{CCH}_2)_2\text{GaCl}\cdot\text{NMe}_3$ . Mol ratio  $(\text{PhMe}_2\text{CCH}_2)_2\text{GaCl}/\text{NMe}_3$ : 1.01. mp 76.5-80.1 °C.  $^1\text{H}$  NMR ( $\text{C}_6\text{H}_6$ ,  $\delta$ ): 1.60 (s, 12H,  $-\text{CMe}_2$ ), 1.52 (s, 9H,  $\text{NMe}_3$ ), 0.86 (s, 4H,  $-\text{CH}_2-$ ).

$(\text{PhMe}_2\text{CCH}_2)\text{GaCl}_2\cdot\text{NMe}_3$ . Mol ratio  $(\text{PhMe}_2\text{CCH}_2)\text{GaCl}_2/\text{NMe}_3$ : 0.92. mp 46.4-48.4 °C.  $^1\text{H}$  NMR ( $\text{C}_6\text{H}_6$ ,  $\delta$ ): 1.67 (s, 15H,  $-\text{CMe}_2$  and  $\text{NMe}_3$ ), 1.57 (s, 2H,  $-\text{CH}_2-$ ).

$(\text{PhMe}_2\text{CCH}_2)_3\text{Ga}\cdot\text{THF}$ . Mol ratio  $\text{Ga}(\text{CH}_2\text{CMe}_2\text{Ph})_3/\text{THF}$ : 0.96. mp 98.7-102.6 °C.  $^1\text{H}$  NMR ( $\text{C}_6\text{H}_6$ ,  $\delta$ ): 3.40 (m, 4H, THF), 1.27 (s, 22H,  $-\text{CMe}_2$  and

THF), 0.80 (s, 6H,  $-\text{CH}_2-$ ).  $(\text{PhMe}_2\text{CCH}_2)_3\text{Ga}\cdot\text{THF}$  was observed to very slowly dissociate at room temperature.

$(\text{PhMe}_2\text{CCH}_2)_2\text{Cl}_2\text{Ga}\cdot\text{THF}$ . Mol ratio  $(\text{PhMe}_2\text{CCH}_2)_2\text{GaCl}_2/\text{THF}$ : 1.05. mp 48.5-49.8 °C.  $^1\text{H}$  NMR ( $\text{C}_6\text{H}_6$ ,  $\delta$ ): 3.34 (m, 4H, THF), 1.49 (s, 6H,  $-\text{CMe}_2$ ), 1.38 (s, 2H,  $-\text{CH}_2-$ ), 0.99 (m, 4H, THF).

### RESULTS AND DISCUSSION

A series of organogallium compounds that incorporate the neophyl ligand including  $\text{Ga}(\text{CH}_2\text{CMe}_2\text{Ph})_3$ ,  $\text{Ga}(\text{CH}_2\text{CMe}_2\text{Ph})_2\text{Cl}$ ,  $\text{Ga}(\text{CH}_2\text{CMe}_2\text{Ph})\text{Cl}_2$  and  $(\text{PhMe}_2\text{CCH}_2)_2\text{GaN}(\text{H})(\text{Pr})$  has been prepared by reactions typical of group 13 chemistry. The homoleptic derivative  $\text{Ga}(\text{CH}_2\text{CMe}_2\text{Ph})_3$  was prepared by a typical Grignard reaction in diethyl ether solution. The ether was readily removed as no apparent adduct formed. However, the isolation of pure samples of  $\text{Ga}(\text{CH}_2\text{CMe}_2\text{Ph})_3$  was hindered by contamination with bineophyl,  $\text{PhMe}_2\text{CCH}_2\text{CH}_2\text{CMe}_2\text{Ph}$ . Fractional distillation and recrystallization served to provide an analytically pure, colorless crystalline solid at room temperature. The halide derivatives were prepared from  $\text{Ga}(\text{CH}_2\text{CMe}_2\text{Ph})_3$  and  $\text{GaCl}_3$  by using stoichiometric ligand redistribution reactions. The neophylgallium amide,  $(\text{PhMe}_2\text{CCH}_2)_2\text{GaN}(\text{H})(\text{Pr})$  was prepared by an elimination reaction between  $\text{Ga}(\text{CH}_2\text{CMe}_2\text{Ph})_3$  and  $\text{H}_2\text{NPr}$  at 140-150 °C.

The characterization data for all compounds, which include partial elemental analyses (C, H), melting points,  $^1\text{H}$  NMR and IR spectroscopic data, cryoscopic molecular weight studies and solubility properties, are consistent with the molecular formulas  $\text{Ga}(\text{CH}_2\text{CMe}_2\text{Ph})_3$ ,  $[\text{Ga}(\text{CH}_2\text{CMe}_2\text{Ph})_2\text{Cl}]_2$ , and  $[\text{Ga}(\text{CH}_2\text{CMe}_2\text{Ph})\text{Cl}_2]_2$ . It is noteworthy that all of the compounds have melting points which are higher than those for the corresponding neopentyl derivatives.<sup>2</sup> The structures of all three compounds would be expected to be

similar to most other organogallium compounds. The homoleptic derivative  $\text{Ga}(\text{CH}_2\text{CMe}_2\text{Ph})_3$  would be expected to be a trigonal planar molecule ( $\text{GaC}_3$ ) whereas the chloride derivatives would be expected to have chloride bridges with four coordinate gallium atoms.

Lewis acid-base studies suggest that the neophylgallium compounds have Lewis acidities which are very similar to those of the corresponding neopentylgallium compounds.<sup>2</sup> If there is a difference, the neophyl derivatives might be slightly stronger Lewis acids. The replacement of a methyl group in the neopentyl group by a phenyl group might serve to reduce steric hindrance slightly and make the neophyl derivatives the stronger Lewis acids. Thus,  $\text{Ga}(\text{CH}_2\text{CMe}_2\text{Ph})_3$  formed a 1:1 adduct with  $\text{NMe}_3$  and THF but the THF adduct appeared to slowly dissociate at room temperature. In contrast, trineopentylgallium did not form an adduct with THF and the characterization data for the  $\text{NMe}_3$  adduct suggested slight dissociation. The monochloride derivative  $\text{Ga}(\text{CH}_2\text{CMe}_2\text{Ph})_2\text{Cl}$  appears to be the weakest Lewis acid of the three derivatives studied as it was the only one of the three new gallium compounds which did not form an adduct with THF. However, an adduct was formed with  $\text{NMe}_3$ . The chloride bridge bonds must compete effectively with the THF for the gallium acid site. The dichloride derivative,  $\text{Ga}(\text{CH}_2\text{CMe}_2\text{Ph})\text{Cl}_2$ , formed 1:1 adducts with both THF and  $\text{NMe}_3$ .

Trineophylgallium(III) undergoes a facile elimination reaction at 140-150 °C with  $\text{NH}_2\text{Pr}$  to form  $(\text{PhMe}_2\text{CCH}_2)_2\text{GaN(H)(Pr)}$  and t-butylbenzene. The gallium-nitrogen product was fully characterized including an X-ray structural study. The unit cell contains one dimeric molecule which lies about a center of symmetry. All molecules are separated by normal van der Waal's distances and there are no anomalously short intermolecular contacts. Distances and angles are provided in Tables 3 and 4.

As shown in Figures 1 and 2, the dimeric molecule contains a  $\text{Ga}_2\text{N}_2$  core which is required by symmetry to be planar. The n-propyl groups are arranged trans to each other across the ring. The Ga-N distances are:  $\text{Ga}(1)-\text{N}(1\text{A}) = \text{Ga}(1\text{A})-\text{N}(1) = 2.013(2)\text{\AA}$  and  $\text{Ga}(1)-\text{N}(1) = \text{Ga}(1\text{A})-\text{N}(1\text{A}) = 2.029(2)\text{\AA}$ . These distances are slightly longer than the Ga-N distances observed for some other related gallium-nitrogen compounds (Table 5). The two gallium atoms are separated by  $2.938(1)\text{\AA}$  whereas the two nitrogen atoms are  $2.776(3)\text{\AA}$  apart. The internal ring angles are:  $\text{N}(1)-\text{Ga}(1)-\text{N}(1\text{A}) = 86.7(1)^\circ$  and  $\text{Ga}(1)-\text{N}(1)-\text{Ga}(1\text{A}) = 93.3(1)^\circ$ . The  $\text{CH}_2\text{CMe}_2\text{Ph}$  ligands are bonded to gallium at distances of  $\text{Ga}(1)-\text{C}(20) = 1.992(2)\text{\AA}$  and  $\text{Ga}(1)-\text{C}(30) = 1.994(3)\text{\AA}$ , with an interligand angle of  $\text{C}(20)-\text{Ga}(1)-\text{C}(30) = 128.3(1)^\circ$ . These ligands are displaced out of the  $\text{Ga}_2\text{N}_2$  plane (as indicated by the  $\text{Ga}(1\text{A})\cdots\text{Ga}-\text{C}$  angles) by  $61.0^\circ$  and  $67.6^\circ$ , respectively. The Ga-C distances are slightly longer than the distances observed for the methyl groups bound to the gallium atoms that form the related  $\text{Ga}_2\text{N}_2$  ring in  $[\text{Me}_2\text{Ga}]_2\text{14aneN}_4[\text{GaMe}_3]_2$ <sup>15</sup> of  $1.933(7)\text{\AA}$  but comparable to the average gallium-carbon distance in  $[(\text{Me}_3\text{CCH}_2)_2\text{GaPPh}_2]_2$ <sup>16</sup> of  $2.003 \pm 0.009\text{\AA}$  and in  $[(t\text{-Bu})_2\text{GaP}(\text{C}_5\text{H}_9)(\text{H})]_2$ <sup>17</sup> of  $2.011(3)\text{\AA}$ . The position of the unique hydrogen atom on nitrogen was refined, yielding an N-H distance of  $0.817(30)\text{\AA}$  and is comparable to the N-H distance in the closely related compound  $[\text{Cl}_2\text{GaN}(\text{H})(\text{SiMe}_3)]_2$ <sup>12</sup> of  $0.838(47)\text{\AA}$ . The unique hydrogen in  $[(\text{PhMe}_2\text{CCH}_2)_2\text{GaN}(\text{H})(\text{Pr})]_2$  is displaced from the gallium-nitrogen plane by  $64.0^\circ$  (defined by the angle  $\text{N}(1\text{A})\cdots\text{N}(1)-\text{H}(1)$ ). The  $\alpha$ -carbon atom of the n-propyl group is displaced from the  $\text{Ga}_2\text{N}_2$  plane by  $43.7^\circ$  and is associated with the distance  $\text{N}(1)-\text{C}(41) = 1.483(3)\text{\AA}$ . Angles around nitrogen are:  $\text{Ga}(1)-\text{N}(1)-\text{C}(41) = 119.3(1)^\circ$ ,  $\text{Ga}(1\text{A})-\text{N}(1)-\text{C}(41) = 120.3(2)^\circ$ ,  $\text{Ga}(1)-\text{N}(1)-\text{H}(1) = 102.5(15)$ ,  $\text{Ga}(1\text{A})-\text{N}(1)-\text{H}(1) = 113.4(15)^\circ$  and  $\text{C}(41)-\text{N}(1)-\text{H}(1) = 106.7(14)^\circ$ .

Symmetry restraints require that the two amide hydrogens are trans to each other across the ring. All other distances and angles in the structure are normal.

The cryoscopic molecular weight studies of  $(\text{PhMe}_2\text{CCH}_2)_2\text{GaN}(\text{H})(\text{Pr})$  identify the presence of dimeric molecules in benzene solution. The  $^1\text{H}$  NMR spectrum of the compound further suggests the presence of only molecules in the trans conformation in solution as only one set of lines for  $\text{CH}_2$  and  $\text{CMe}_2$  protons for the neophyl ligand or for the propyl group protons was observed. If the cis isomer had been present, two sets of lines of equal intensity, one for each type of proton, would have been expected. Thus, the molecule in solution is the same as the molecule present in the crystalline solid. There was no change in either the degree of association or in the conformation of the molecule upon dissolution in benzene. In comparison,  $[\text{Me}_2\text{GaN}(\text{Me})(\text{Ph})]_2^{18}$  existed as a mixture of cis (70%) and trans (30%) isomers in benzene solution. The sharp melting point of  $[\text{Me}_2\text{GaN}(\text{Me})(\text{Ph})]_2$  (112-114 °C) and the presence of only one crystalline form was used to suggest only one isomer (probably trans) in the solid state. The related compound  $[\text{Me}_2\text{InN}(\text{Me})(\text{Ph})]_2^{18}$  existed as a trans isomer in the solid state according to a structural study whereas a mixture of cis and trans isomers were observed in solution. Thus, at least one metal nitrogen bond in these dimeric species was broken upon dissolution in order to have isomerization.

Acknowledgment. This work was supported in part by the Office of Naval Research and by a generous grant from Eastman Kodak Co. We thank the National Science Foundation for providing funds for the Siemens R3m/V diffractometer through a grant from the Chemical Instrumentation Program (89-13733).



Supplementary Material. Complete lists of distances and angles, anisotropic thermal parameters, calculated positions of hydrogen atoms and  $F_o/F_c$  list for  $[(\text{PhMe}_2\text{CCH}_2)_2\text{Ga}(\text{H})(\text{Pr})]_2$  (    pages). For ordering information, see any current masthead page.


# References

1. Beachley, O. T., Jr.; Simmons, R. G. Inorg. Chem. 1980, 19, 1021.
2. Beachley, O. T., Jr.; Pazik, J. C. Organometallics 1988, 7, 1516.
3. Beachley, O. T., Jr.; Churchill, M. R.; Pazik, J. C.; Ziller, J. W. Organometallics 1986, 5, 1814.
4. Beachley, O. T., Jr.; Churchill, M. R.; Pazik, J. C.; Ziller, J. W. Organometallics 1987, 6, 2088.
5. Beachley, O. T., Jr.; Hallock, R. B. Organometallics 1987, 6, 170.
6. Beachley, O. T., Jr.; Hallock, R. B.; Zhang, H. M.; Atwood, J. L. Organometallics 1985, 4, 1675.
7. Shriver, D. F.; Drezzdon, M. A. "The Manipulation of Air-Sensitive Compounds; Wiley: New York, 1986, p 38.
8. Whitesides, G. M.; Panek, E. J.; Stedronsky, E. R. J. Am. Chem. Soc. 1972, 94, 232.
9. Richards, D. H.; Scilly, N. F. J. Chem. Soc. (C) 1969, 55.
10. Siemens SHELXTL PLUS Manual, 2nd Edition (1990). Siemens Analytical Instruments, Madison, Wisconsin.
11. Churchill, M. R. Inorg. Chem. 1973, 12, 1213.
12. Nutt, W. R.; Anderson, J. A.; Odom, J. D.; Williamson, M. M.; Rubin, B. H. Inorg. Chem. 1985, 24, 159.
13. Harrison, W.; Storr, A.; Trotter, J. J. Chem. Soc. Chem. Comm. 1971, 1101.
14. Hwang, J.-W.; Hanson, S. A.; Britton, D.; Evans, J. F.; Jensen, K. F.; Gladfelter, W. L. Chem. Mat. 1990, 2, 342.
15. Lee, B.; Pennington, W. T.; Robinson, G. H.; Rogers, R. D. J. Organomet. Chem. 1990, 396, 269.

16. Banks, M. A.; Beachley, O. T., Jr.; Buttrey, L. A.; Churchill, M. R.;  
Fettingner, J. C. Organometallics 1991, 10, 1901.
17. Heaton, D. E.; Jones, R. A.; Kidd, K. B.; Cowley, A. H.; Nunn, C. M.  
Polyhedron 1988, 7, 1901.
18. Beachley, O. T., Jr.; Bueno, C.; Churchill, M. R.; Hallock, R. B.;  
Simmons, R. G. Inorg. Chem. 1981, 20, 2423.

Table 1.

Experimental Data for the X-Ray Diffraction Study of trans-



Crystal Data

|                        |   |
|------------------------|---|
| Empirical Formula      | $\text{C}_{46} \text{H}_{68} \text{Ga}_2 \text{N}_2$  |
| Color; Habit           | White   |
| Crystal Size (mm)      | .35x.30x.30   |
| Crystal System         | Triclinic   |
| Space Group            | $\text{P}\bar{1}$   |
| Unit Cell Dimensions   | $\underline{a} = 10.739(2) \text{ \AA}$<br>$\underline{b} = 10.844(2) \text{ \AA}$<br>$\underline{c} = 11.125(2) \text{ \AA}$<br>$\alpha = 69.140(10)^\circ$<br>$\beta = 62.960(10)^\circ$<br>$\gamma = 76.740(10)^\circ$ |
| Volume                 | $1074.7(3) \text{ \AA}^3$   |
| Z                      | 1   |
| Formula weight         | 788.5   |
| Density(calc.)         | $1.218 \text{ Mg/m}^3$  |
| Absorption Coefficient | $1.279 \text{ mm}^{-1}$   |
| F(000)                 | 420   |

Data Collection

|                         |   |
|-------------------------|---|
| Diffractometer Used     | Siemens R3m/V   |
| Radiation               | MoK $\alpha$ ( $\lambda$ = 0.71073 Å)   |
| Temperature (K)         | 298   |
| Monochromator           | Highly oriented graphite crystal  |
| 2 $\theta$ Range        | 5.0 to 50.0°  |
| Scan Type               | 2 $\theta$ - $\theta$   |
| Scan Speed              | Constant; 1.00°/min. in $\omega$  |
| Scan Range ( $\omega$ ) | 0.70° plus K $\alpha$ -separation   |
| Background Measurement  | Stationary crystal and stationary counter at beginning and end of scan, each for 25.0% of total scan time |
| Index Ranges            | $-12 \leq h \leq 12$ , $-12 \leq k \leq 12$<br>$-13 \leq l \leq 13$                                       |
| Reflections Collected   | 7606  |
| Independent Reflections | 3803 ( $R_{\text{int}}$ = 1.93%)  |
| Observed Reflections    | 3803 ( $F > 0.3\sigma(F)$ )   |
| Observed Reflections    | 3111 ( $F > 6.0\sigma(f)$ )   |
| Absorption Correction   | Semi-empirical  |
| Min./Max. Transmission  | 0.8180 / 0.9835   |

Solution and Refinement

|                                      |  |
|--------------------------------------|--|
| System Used                          | Siemens SHELXTL PLUS (VMS)   |
| Solution                             | Direct Methods   |
| Refinement Method                    | Full-Matrix Least-Squares  |
| Quantity Minimized                   | $\sum w(F_o - F_c)^2$  |
| Absolute Structure                   | N/A  |
| Extinction Correction                | $\chi = 0.0004(2)$ , where<br>$F^* = F [ 1 + 0.002\chi F^2 / \sin(2\theta) ]^{-1/4}$ |
| Hydrogen Atoms                       | Riding model, fixed isotropic U  |
| Weighting Scheme                     | $w^{-1} = \sigma^2(F) + 0.0005F^2$   |
| Number of Parameters refined         | 264  |
| Final R indices (0.25 $\sigma$ data) | R = 3.57 %, wR = 3.46 %  |
| Final R indices (6.00 $\sigma$ data) | R = 2.52 %, wR = 2.62 %  |
| R Indices (all data)                 | R = 3.57 %, wR = 3.46 %  |
| Goodness-of-Fit                      | 0.97   |
| Largest and Mean $\Delta/\sigma$     | 0.002, 0.000   |
| Data-to-Parameter Ratio              | 14.4:1   |
| Largest Difference Peak              | 0.30 eÅ <sup>-3</sup>  |
| Largest Difference Hole              | -0.22 eÅ <sup>-3</sup>   |

Table 2.

Final Atomic Coordinates ( $\times 10^4$ ) and Equivalent Isotropic Displacement Coefficients ( $\text{\AA} \times 10^3$ ) for trans- $[(\text{PhMe}_2\text{CCH}_2)_2\text{GaN(H)(Pr)}]_2$ .

|       | x        | y        | z        | U(eq)  |
|-------|----------|----------|----------|--------|
| Ga(1) | 3757(1)  | 4258(1)  | 5639(1)  | 30(1)  |
| N(1)  | 5819(2)  | 4025(2)  | 4368(2)  | 32(1)  |
| H(1)  | 5794(23) | 4060(21) | 3633(23) | 33(6)  |
| C(20) | 3404(2)  | 2824(2)  | 7450(2)  | 43(1)  |
| C(21) | 2207(3)  | 2956(2)  | 8868(2)  | 45(1)  |
| C(22) | 2117(3)  | 1598(2)  | 9989(2)  | 47(1)  |
| C(23) | 3115(3)  | 1106(3)  | 10522(3) | 84(2)  |
| C(24) | 3076(4)  | -123(4)  | 11489(4) | 99(2)  |
| C(25) | 2049(4)  | -907(3)  | 11932(3) | 86(2)  |
| C(26) | 1034(4)  | -451(3)  | 11444(3) | 80(2)  |
| C(27) | 1068(3)  | 792(3)   | 10480(3) | 63(1)  |
| C(28) | 2478(4)  | 3991(3)  | 9329(3)  | 76(2)  |
| C(29) | 819(3)   | 3416(3)  | 8687(3)  | 66(1)  |
| C(30) | 2733(2)  | 4745(2)  | 4402(2)  | 41(1)  |
| C(31) | 2441(2)  | 3672(2)  | 3966(2)  | 46(1)  |
| C(32) | 3808(3)  | 2933(2)  | 3200(2)  | 46(1)  |
| C(33) | 4793(3)  | 3586(3)  | 1917(3)  | 61(1)  |
| C(34) | 6015(4)  | 2928(4)  | 1196(3)  | 84(2)  |
| C(35) | 6291(4)  | 1608(4)  | 1733(5)  | 94(2)  |
| C(36) | 5347(5)  | 951(4)   | 3006(5)  | 101(3) |
| C(37) | 4123(4)  | 1602(3)  | 3717(4)  | 75(2)  |
| C(38) | 1439(3)  | 2722(3)  | 5289(3)  | 64(1)  |
| C(39) | 1681(3)  | 4317(3)  | 2986(3)  | 70(2)  |
| C(41) | 6649(2)  | 2774(2)  | 4756(2)  | 41(1)  |
| C(42) | 8130(3)  | 2664(2)  | 3672(3)  | 50(1)  |
| C(43) | 8934(3)  | 1364(3)  | 4087(3)  | 72(2)  |

\* Equivalent isotropic U defined as one third of the trace of the orthogonalized  $U_{ij}$  tensor

Table 3

Important Interatomic Distances (Å) for trans-[(PhMe<sub>2</sub>CCH<sub>2</sub>)<sub>2</sub>GaN(H)(Pr)]<sub>2</sub>.

|                |          |             |           |
|----------------|----------|-------------|-----------|
| Ga(1)-N(1)     | 2.029(2) | Ga(1)-C(20) | 1.992(2)  |
| Ga(1)-N(1A)    | 2.013(2) | Ga(1)-C(30) | 1.994(3)  |
| Ga(1)···Ga(1A) | 2.938(1) | N(1)-C(41)  | 1.483(3)  |
| N(1)···N(1A)   | 2.776(3) | N(1)-H(1)   | 0.817(30) |



Table 4

Important Interatomic Angles (°) for trans-[(PhMe<sub>2</sub>CCH<sub>2</sub>)<sub>2</sub>GaN(H)(Pr)]<sub>2</sub>.

|                   |           |                   |           |
|-------------------|-----------|-------------------|-----------|
| N(1)-Ga(1)-N(1A)  | 86.7(1)   | C(20)-Ga(1)-C(30) | 128.3(1)  |
| N(1)-Ga(1)-C(20)  | 107.1(1)  | N(1A)-Ga(1)-C(20) | 114.3(1)  |
| N(1)-Ga(1)-C(30)  | 105.9(1)  | N(1A)-Ga(1)-C(30) | 106.2(1)  |
| Ga(1)-N(1)-Ga(1A) | 93.3(1)   | Ga(1)-N(1)-C(41)  | 119.3(1)  |
| Ga(1)-N(1)-H(1)   | 102.5(15) | Ga(1A)-N(1)-C(41) | 120.3(2)  |
| Ga(1A)-N(1)-H(1)  | 113.4(15) | C(41)-N(1)-H(1)   | 106.4(14) |

Table 5.

Comparisons of Ga-N and GaC Bond Distances in Some Gallium-Nitrogen Compounds.

| <u>Compound</u>   | <u>Ga-N</u> | <u>Ga-C</u> | <u>Reference</u> |
|---|-------------|-------------|------------------|
| $[\text{Cl}_2\text{GaN}(\text{H})(\text{SiMe}_3)]_2$                  | 1.974(4)    | --          | 12               |
|   | 1.964(4)    | --          |                  |
| $[\text{Cl}_2\text{GaN}(\text{Me})(\text{SiMe}_3)]_2$                 | 1.985(5)    | --          | 12               |
|   | 1.983(5)    | --          |                  |
|   | 1.987(5)    | --          |                  |
|   | 1.989(5)    | --          |                  |
| $[\text{H}_2\text{GaN}(\text{CH}_2)_2]_3$                             | 1.97(2)     | --          | 13               |
| $[\text{H}_2\text{GaN}(\text{H})_2]_3$                                | 1.96(2)     | --          | 14               |
|   | 1.96(2)     | --          |                  |
|   | 1.98(2)     | --          |                  |
|   | 1.96(1)     | --          |                  |
| $[\text{Me}_2\text{Ga}]_2[14]\text{ane } \text{N}_4[\text{GaMe}_3]_2$ | 2.014(6)    | 1.933(7)    | 15               |
|   | 2.003(6)    |             |                  |
| $[(\text{PhMe}_2\text{CCH}_2)_2\text{GaN}(\text{H})(\text{Pr})]_2$    | 2.013(2)    | 1.992(2)    | This work        |
|   | 2.029(2)    | 1.994(3)    |                  |

Figure 1. The  $[(\text{PhMe}_2\text{CCH}_2)\text{GaN}(\text{H})(\text{Pr})]_2$  molecule, with all C-bonded hydrogen atoms omitted.

Figure 2. The  $[\text{Ga}(\text{C})_2(\mu\text{-NHC}_3)]_2$  core of the molecule.




FIG 1




FIG. 2

TECHNICAL REPORT DISTRIBUTION LIST - GENERAL

Office of Naval Research (2)\*  
Chemistry Division, Code 1113  
800 North Quincy Street  
Arlington, Virginia 22217-5000

Dr. Richard W. Drisko (1)  
Naval Civil Engineering  
Laboratory  
Code L52  
Port Hueneme, CA 93043

Dr. James S. Murday (1)  
Chemistry Division, Code 6100  
Naval Research Laboratory  
Washington, D.C. 20375-5000

Dr. Harold H. Singerman (1)  
David Taylor Research Center  
Code 283  
Annapolis, MD 21402-5067

Dr. Robert Green, Director (1)  
Chemistry Division, Code 385  
Naval Weapons Center  
China Lake, CA 93555-6001

Chief of Naval Research (1)  
Special Assistant for Marine  
Corps Matters  
Code 00MC  
800 North Quincy Street  
Arlington, VA 22217-5000

Dr. Eugene C. Fischer (1)  
Code 2840  
David Taylor Research Center  
Annapolis, MD 21402-5067

Defense Technical Information  
Center (2)  
Building 5, Cameron Station  
Alexandria, VA 22314

Dr. Elek Lindner (1)  
Naval Ocean Systems Center  
Code 52  
San Diego, CA 92152-5000

Commanding Officer (1)  
Naval Weapons Support Center  
Dr. Bernard E. Doua  
Crane, Indiana 47522-5050

\* Number of copies to forward